arXiv:1112.1706v2 [hep-th] 11 Dec 2011

On generalized analogues of black brane solutions in the model with multicomponent anisotropic fluid

V.D. Ivashchuk[1]

Center for Gravitation and Fundamental Metrology, VNIIMS, 3/1 M. Ulyanovoy Str., Moscow 119313, Russia and
Institute of Gravitation and Cosmology, PFUR, 6 Miklukho-Maklaya Str., Moscow 117198, Russia

Abstract

A family of spherically symmetric solutions in the model with m-component anisotropic fluid is considered. The metrics are defined on a manifold which contains a product of $n-1$ Ricci-flat "internal" spaces. The equation of state for any s-th component is defined by a vector $U^s = (U^s_i)$ belonging to $\mathbb{R}^{n+1}$ and obeying inequalities $U^s_1 = q_s > 0$, $s = 1, \ldots, m$. The solutions are governed by moduli functions H_s which are solutions of (master) non-linear differential equations with certain boundary conditions imposed. It is shown that for coinciding $q_s = q$ there exists a subclass of solutions with regular horizon when $q = 1, 2, \ldots$ and U^s-vectors correspond to certain semisimple Lie algebras. An extension of these solutions to block-orthogonal set of vectors U^s with natural parameters q_s coinciding inside blocks is also proposed. Generalized q-analogues of black brane solutions are suggested and q-analogue of $M_2 - M_5$ dyonic solution in $D = 11$ supergravity (corresponding to Lie algebra A_2) is considered.

PACS number(s): 04.50.+h, 04.20.Jb, 04.70.Bw

Key words: black branes, multicomponent anisotropic fluid, Lie algebras, polynomials

---

[1] ivashchuk@mail.ru

1 Introduction

In this paper we consider spherically-symmetric solutions with a horizon in the multidimensional model with multicomponent anisotropic fluid defined on warped product manifolds $\mathbb{R} \times M_0 \times \ldots \times M_n$. Here we generalize solutions from [1]-[5] which may simulate black brane solutions appearing in the models with antisymmetric forms and scalar fields. (For black brane solutions see [6]-[27] and references therein.)

Here we present a new family of exact spherically-symmetric solutions in the model with m-component anisotropic fluid for the following equations of state [4]

$$p_r^s = -\rho^s/(2q_s - 1), \qquad p_0^s = \rho^s/(2q_s - 1), \tag{1.1}$$

and

$$p_i^s = \left(1 - \frac{2U_i^s}{d_i}\right) \rho^s/(2q_s - 1), \tag{1.2}$$

$i > 1$, $s = 1, \ldots, m$, where for s-th component: ρ^s is a density, p_r^s is a radial pressure, p_i^s is a pressure in M_i ($i > 1$) and $q_s \neq 1/2$. Parameters U_i^s ($i > 1$) and $q_s = U_1^s > 0$ obey certain relations described in Section 2. Here the manifold M_0 is d_0-dimensional sphere and p_0^s is the pressure in the tangent direction. (For multifactor cosmological models see [28, 29, 30, 31] and refs. therein.)

Special solutions with $q_s = 1$ were found in [1] (for $m = 1$), [2] (for orthogonal vectors $U^s = (U_i^s)$) and [5]. For $q_s > 0$ special solutions were obtained in [4] (for orthogonal U^s) and in [3] (for $m = 1$). In [27] the post-Newtonian parameters β and γ corresponding to the 4-dimensional section of the metric from [5] were calculated.

Here we show that there exist solutions with regular horizon when U^s-vectors correspond to simple roots of certain semisimple Lie algebra and q_s are integer numbers obeying certain restrictions. (For simple Lie algebra q_s should coincide: $q_s = q$ for all s.) We construct generalized analogues of black brane solutions. This may be of interest for a possible search of (holographically) dual quantum models.

The paper is organized as follows. In Section 2 the model with multicomponent anisotropic (or "perfect") fluid is formulated. In Section 3 a subclass of spherically symmetric solutions (generalizing those from [5]) is presented.

In Section 4 and a subclass of solutions with regular horizon corresponding to integer q_s is singled out. In Section 5 q-analogs of black brane solutions are considered, e.g. a q-analogue of $M2 \cap M5$ dyonic solution.

2 The model

Here, we consider a family of spherically symmetric solutions to Hilbert-Einstein equations with an multicomponent anisotropic fluid matter source

$$R^M_N - \frac{1}{2}\delta^M_N R = kT^M_N \tag{2.1}$$

defined on the manifold

$$M = \mathbb{R}_{.} \times (M_0 = S^{d_0}) \times (M_1 = \mathbb{R}) \times M_2 \times \ldots \times M_n \tag{2.2}$$

with the block-diagonal metrics

$$g = e^{2\gamma(u)} du \otimes du + \sum_{i=0}^{n} e^{2\beta^i(u)} h^{[i]}. \tag{2.3}$$

Here $\mathbb{R}_{.} = (u_-, u_+)$ is an interval. The manifold M_i with the metric $h^{[i]}$ is the Ricci-flat space of dimension d_i, $i = 1, 2, \ldots, n$ and $h^{(0)}$ is standard metric on the unit sphere S^{d_0} ($\mathrm{Ric}[h^{[0]}] = (d_0 - 1)h^{[0]}$), u is radial variable, κ is the multidimensional gravitational constant, $d_1 = 1$ and $h^{[1]} = -dt \otimes dt$.

The energy-momentum tensor is adopted in the following form

$$T^M_N = \sum_{s=1}^{m} T^{(s)M}_N, \tag{2.4}$$

where

$$T^{(s)M}_N = \mathrm{diag}(p^s_r, p^s_0 \delta^{m_0}_{k_0}, -\rho^s, p^s_2 \delta^{m_2}_{k_2}, \ldots, p^s_n \delta^{m_n}_{k_n}), \tag{2.5}$$

$q_s > 0$ and $q_s \neq 1/2$. The pressures $p^s_r = p^s_r(u)$, $p^s_i = p^s_i(u)$ and the densities $\rho^s = \rho^s(u)$ obey the relations (1.1), (1.2).

As in [29, 31] we impose the "conservation law" restrictions

$$\nabla_M T^{(s)M}_N = 0, \tag{2.6}$$

$s = 1, \ldots, m - 1$. In what follows we put $\kappa = 1$ for simplicity.

As it was mentioned in [4] it is more convenient for finding of exact solutions, to write the stress-energy tensor in cosmological-type form

$$(T_N^{(s)M}) = \mathrm{diag}(-\hat{\rho}^s, \hat{p}_0^s \delta_{k_0}^{m_0}, \hat{p}_1^s \delta_{k_1}^{m_1}, \ldots, \hat{p}_n^s \delta_{k_n}^{m_n}), \tag{2.7}$$

where $\hat{\rho}^s$ and $\hat{p}_i^s$ are "effective" density and pressures of s-th component, respectively, depending upon u. The physical density ρ^s and pressures p_i^s are related to the effective ones by formulas

$$\rho^s = -\hat{p}_1^s, \quad p_r^s = -\hat{\rho}^s, \quad p_i^s = \hat{p}_i^s \quad (i \neq 1), \tag{2.8}$$

$s = 1, \ldots, m$.

The equations of state may be written in the following form

$$\hat{p}_i = \left(1 - \frac{2U_i^s}{d_i}\right) \hat{\rho}^s, \tag{2.9}$$

where U_i^s are constants, $i = 0, 1, \ldots, n$.

We impose the following restrictions on $U^s = (U_i^s) \in \mathbb{R}^{n+1}$:

$$1^o. \quad U_0^s = 0, \tag{2.10}$$

$$2^o. \quad U_1^s = q_s > 0, \tag{2.11}$$

$$3^o. \quad ((U^s, U^l)) = (U_i^s G^{ij} U_j^l) \quad \text{is non} - \text{degenerate matrix.} \tag{2.12}$$

Here

$$G^{ij} = \frac{\delta^{ij}}{d_i} + \frac{1}{2 - D} \tag{2.13}$$

are components of the matrix inverse to the matrix of the minisuperspace metric [28]

$$(G_{ij}) = (d_i \delta_{ij} - d_i d_j), \tag{2.14}$$

$i, j = 0, \ldots, n$, and $D = 1 + \sum\limits_{i=0}^{n} d_i$ is the total dimension.

In our case the scalar products (2.12) are given by relations:

$$(U^s, U^l) = \sum_{i=1}^{n} \frac{U_i^s U_i^l}{d_i} + \frac{1}{2 - D} \left(\sum_{i=1}^{n} U_i^s\right) \left(\sum_{j=1}^{n} U_j^l\right). \tag{2.15}$$

It was proved in Appendix of [4] that the relations 1^o and 2^o imply $K_s = (U^s, U^s) > 0$ for all s. In what follows we will use a "quasi-Cartan" matrix

$$2(U^s, U^l)/(U^l, U^l) = A_{sl}, \tag{2.16}$$

which is a non-degenerate matrix due to (2.12).

3 Exact solutions

For the equations of state (1.1) and (1.2) with parameters U_i^s $(i > 0)$ obeying (2.12) (see (2.15)) we obtain the following spherically symmetric solutions to the Hilbert-Einstein equations (2.1)

$$g = J_0 \left(F^{-1} dR \otimes dR + R^2 h^{[0]} \right) - J_1 F dt \otimes dt + \sum_{i=2}^{n} J_i h^{[i]}, \tag{3.1}$$

$$\rho^s = -(2q_s - 1) \frac{A_s F^{q_s - 1}}{J_0 R^{2d_0}} \prod_{l=1}^{m} H_l^{-A_{sl}}, \tag{3.2}$$

which may be derived by analogy with the black brane solutions [22, 23]. Here $F = 1 - 2\mu R^{-d}$, $d = d_0 - 1$, $h^{[0]}$ is canonical metric on S^{d_0} and Ric$[h^{[i]}]$= 0 for $i > 0$,

$$J_i = \prod_{s=1}^{m} H_s^{-2h_s U^{si}}, \tag{3.3}$$

$i = 0, 1, ..., n$; $\mu > 0$ is an integration constant and

$$U^{si} = G^{ij} U_j^s = \frac{U_i^s}{d_i} + \frac{1}{2 - D} \sum_{j=0}^{n} U_j^s, \tag{3.4}$$

$$h_s = K_s^{-1}, \quad K_s = (U^s, U^s). \tag{3.5}$$

It follows from 1^o and (3.4) that

$$U^{s0} = \frac{1}{2 - D} \sum_{j=1}^{n} U_j^s. \tag{3.6}$$

Functions $H_s > 0$ obey the equations

$$R^{d_0} \frac{d}{dR} \left[R^{d_0} \frac{F}{H_s} \frac{dH_s}{dR} \right] = B_s F^{q_s - 1} \prod_{l=1}^{m} H_l^{-A_{sl}}, \tag{3.7}$$

with $B_s = 2K_s A_s$ and the boundary conditions imposed:

$$H_s \to H_{s0} \neq 0 \quad \text{for} \quad R^d \to 2\mu, \tag{3.8}$$

and

$$H_s(R = +\infty) = 1, \tag{3.9}$$

$s = 1, ..., m$.

The first boundary condition (3.8) implies the fulfillment of the weak horizon condition (i.e. infinite time of propagation of light for $R^d \to 2\mu$). The second one (3.9) leads to an asymptotically flatness of the $(2 + d_0)$-dimensional section of the metric.

Due to to (2.10) and (3.4) the metric may be rewritten as follows

$$g = J_0 \Biggl[F^{-1} dR \otimes dR + R^2 h^{[0]} - \left(\prod_{s=1}^{m} H_s^{-2q_s h_s} \right) F dt \otimes dt \tag{3.10}$$
$$+ \sum_{i=2}^{n} Y_i h^{[i]} \Biggr],$$

where

$$Y_i = \prod_{s=1}^{m} H_s^{-2h_s U_i^s / d_i}. \tag{3.11}$$

The solution (3.1), (3.2) may be verified just by a substitution into equations of motion. (A complete derivation of this solution will be given in a separate paper [33].)

Remark. The solution is valid for arbitrary Einstein space $(M_0, h^{[0]})$ obeying $\mathrm{Ric}[h^{[0]}] = (d_0 - 1) h^{[0]}$.

A special orthogonal case of this solution (when $(U^s, U^l) = 0$ for $s \neq l$) was considered earlier in [4] (for $m = 1$ see [3]). The solution with $q_s = 1$ was considered in [5]; for special orthogonal case see also [2] and [1] (for $m = 1$).

4 Solutions with a regular horizon.

Here we consider a subclass of black brane like solutions with a regular horizon. We impose the following (additional) condition on the solutions

$$H_s(R) > 0 \text{ is smooth in } (R_\epsilon, +\infty), \tag{4.1}$$

$s = 1, ..., m$, where $R_\epsilon = (2\mu)^{1/d} e^{-\epsilon}$ with some $\epsilon > 0$. Then the metric (3.1) has a regular horizon at $R^d = 2\mu$.

4.1 One-block solutions.

Let us put

$$q_s = q \tag{4.2}$$

for all s, where q is a natural number: $q = 1, 2, \ldots$. In this case there exists a subclass of solutions with regular horizon at $R^d = 2\mu$.

Let us denote $z = R^{-d} \in (0, (2\mu)^{-1})$ and

$$2\mu Z = 1 - (1 - 2\mu z)^q \tag{4.3}$$

where $Z \in (0, (2\mu)^{-1})$ $(q > 0)$.

We get a set of equations

$$\frac{d}{dZ}\left(\frac{(1 - 2\mu Z)}{H_s}\frac{d}{dZ}H_s\right) = \bar{B}_s \prod_{l=1}^{m} H_l^{-A_{sl}}, \tag{4.4}$$

where $H_s = H_s(Z) > 0$ for $Z \in (0, (2\mu)^{-1})$ and $\bar{B}_s = B_s/(qd)^2 \neq 0$. Eqs. (3.8) and (3.9) read

$$H_s((2\mu)^{-1} - 0) = H_{s0} \in (0, +\infty), \tag{4.5}$$

$$H_s(+0) = 1, \tag{4.6}$$

$s = 1, ..., m$.

The condition (4.1) reads as follows

$$H_s(Z) > 0 \text{ is smooth in } (0, Z_\epsilon), \tag{4.7}$$

$s = 1, ..., m$, where $Z_\epsilon = (2\mu)^{-1} e^{\epsilon d}$ with some $\epsilon > 0$.

It was conjectured in [21] that equations (4.4)-(4.6) have polynomial solutions when $(A_{ss'})$ is a Cartan matrix for some semisimple finite-dimensional Lie algebra $\mathbf{G}$ of rank m. In this case we get

$$H_s(Z) = 1 + \sum_{k=1}^{n_s} P_s^{(k)} Z^k, \tag{4.8}$$

where $P_s^{(k)}$ are constants, $k = 1, \ldots, n_s$; $P_s^{(n_s)} \neq 0$, and

$$n_s = b_s \equiv 2 \sum_{l=1}^{m} A^{sl} \tag{4.9}$$

$s = 1, ..., m$, are the components of twice the dual Weyl vector in the basis of simple co-roots [32]. Here $(A^{sl}) = (A^{-1}_{sl})$.

This conjecture was verified for $\mathbf{A_m}$ and $\mathbf{C_{m+1}}$ series of Lie algebras in [22, 23]. In the extremal case ($\mu = +0$) an analogue of this conjecture was suggested (implicitly) in [15].

$\mathbf{A_1}$ -case. The simplest example occurs for $m = 1$ when $(A_{sl}) = (2)$ is a Cartan matrix for the simple Lie algebra $\mathbf{A_1}$. In this case

$$H_1(Z) = 1 + P_1 Z, \tag{4.10}$$

with $P_1 \neq 0$, satisfying

$$P_1(P_1 + 2\mu) = -\bar{B}_1 = -2K_1 A_1/(qd)^2, \tag{4.11}$$

$s = 1, ..., m$. When $A_1 < 0$ (or, equivalently, $\rho^1 > 0$) there exists a unique number $P_1 > 0$ obeying (4.11). In this case we are led to a solution from [3].

A_2-case. For the Lie algebra $\mathcal{G}$ coinciding with $\mathbf{A_2} = sl(3)$ we get $n_1 = n_2 = 2$ and

$$H_s = 1 + P_s Z + P^{(2)}_s Z^2, \tag{4.12}$$

where $P_s = P^{(1)}_s$ and $P^{(2)}_s \neq 0$ are constants, $s = 1, 2$.

It was found in [21] that for $P_1 + P_2 + 4\mu \neq 0$ (e.g. when all $P_s > 0$) the following relations take place

$$P^{(2)}_s = \frac{P_s P_{s+1}(P_s + 2\mu)}{2(P_1 + P_2 + 4\mu)}, \qquad \bar{B}_s = -\frac{P_s(P_s + 2\mu)(P_s + 4\mu)}{P_1 + P_2 + 4\mu}, \tag{4.13}$$

$s = 1, 2$. Here we denote $s + 1 = 2, 1$ for $s = 1, 2$, respectively.

"Master" equations (4.4) were integrated in [25, 26] for Lie algebras $\mathbf{C_2}$ and $\mathbf{A_3}$, respectively. (For $\mathbf{D_4}$-solutions in the extremal case $\mu \to +0$ see [15].)

4.2 Block-orthogonal solutions.

Let $S = \{1, \ldots, m\}$ and

$$S = S_1 \cup \ldots \cup S_k, \qquad S_a \cap S_b = \emptyset, \quad a \neq b, \tag{4.14}$$

$S_a \neq \emptyset$, i.e. the set S is a union of k non-intersecting (non-empty) subsets $S_1, \dots, S_k$. Here $S_1 < \dots < S_k$ and the ordering in S_i is inherited by the ordering in S.

We put

$$(U^s, U^l) = 0, \tag{4.15}$$

for all $s \in S_a$, $l \in S_b$, $a \neq b$; $a, b = 1, \dots, k$.

Let q_s be coinciding inside blocks, i. e.

$$q_s = q_{(a)}, \tag{4.16}$$

for all $s \in S_a$, where $q_{(a)}$ is a natural number:

$$q_{(a)} = 1, 2, \dots \tag{4.17}$$

for any $a = 1, \dots, k$.

Master equations (3.7) in this case split into k sets of equations

$$\frac{d}{dZ_a}\left(\frac{(1 - 2\mu Z_a)}{H_s}\frac{d}{dZ_a}H_s\right) = \bar{B}_s \prod_{l \in S_a} H_l^{-A_{sl}}, \tag{4.18}$$

$s \in S_a$, $\bar{B}_s = B_s/(q_{(a)}d)^2 \neq 0$, and

$$2\mu Z_a = 1 - (1 - 2\mu z)^{q_{(a)}}, \tag{4.19}$$

$a = 1, \dots, k$.

$\mathbf{A_1} \oplus \dots \oplus \mathbf{A_1}$ -case. A simple example occurs in the orthogonal case: $(U^s, U^l) = 0$ for $s \neq l$ [14, 16]. In this case $(A_{sl}) = \mathrm{diag}(2, \dots, 2)$ is a Cartan matrix for the semisimple Lie algebra $\mathbf{A_1} \oplus \dots \oplus \mathbf{A_1}$ and

$$H_s = 1 + P_s Z_s, \tag{4.20}$$

with $P_s \neq 0$, satisfying

$$P_s(P_s + 2\mu) = -\bar{B}_s = -2K_s A_s/(q_s d)^2, \tag{4.21}$$

$s = 1, ..., m$. When all $A_s < 0$ (or, equivalently, $\rho^s > 0$) there exists a unique set of numbers $P_s > 0$ obeying (4.21). Here we are led to a solution from [4].

Analogously one can consider a case of semisimple Lie algebra which is a sum of k simple Lie algebras: $\mathbf{G} = \mathbf{G_1} \oplus \dots \oplus \mathbf{G_k}$. In this case a natural number $q_{(a)}$ should be assigned to a-th component $\mathbf{G_a}$, $a = 1, \dots, k$.

There exists also a special solution to eqs. (4.18) [17, 18, 19]

$$H_s(Z_a) = (1 + P_s Z_a)^{b_s} \tag{4.22}$$

with b_s from (4.9) and parameters P_s coinciding inside a-th block , i.e. $P_s = P_{s'}$ for $s, s' \in S_a$. Parameters $P_s \neq 0$ satisfy the relations

$$P_s(P_s + 2\mu) = -\bar{B}_s/b_s, \tag{4.23}$$

$b_s \neq 0$ and parameters $\bar{B}_s/b^s$ are also coinciding inside blocks, i.e. $\bar{B}_s/b_s = \bar{B}_{s'}/b_{s'}$ for $s, s' \in S_a$. Here $a = 1, \ldots, k$.

5 q-analogs of black brane solutions

The solutions with a regular horizon from [5] allowed us to simulate certain intersecting black brane solutions [24] in the model with antisymmetric forms (without scalar fields) when all $q_s = 1$.

These solutions may be also generalized to the block-ortogonal case (4.14)-(4.17) for a chosen set of natural $q_{(a)} \in \mathbb{N}$, $a = 1, ..., k$.

We put

$$U_i^s = q_{(a)} d_i, \quad i \in I_s; \qquad U_i^s = 0, \quad i \notin I_s, \tag{5.1}$$

for all $s \in S_a$, $a = 1, ..., k$.

Here $I_s = \{i_1, \ldots, i_k\} \in \{1, \ldots n\}$ is the index set [24] corresponding to brane submanifold $M_{i_1} \times \ldots \times M_{i_k}$. All sets I_s contain index 1, corresponding to the time manifold $M_1 = \mathbb{R}$.

The relation (5.1) leads us to the following relation for dimensions of intersections of brane submanifolds ("worldvolumes") [16, 24]:

$$q_{(a)} q_{(b)} \left[d(I_s \cap I_l) - \frac{d(I_s) d(I_l)}{D - 2} \right] = (U^s, U^l), \tag{5.2}$$

$s \in S_a$, $l \in S_b$; $a, b = 1, ..., k$. Here $d(I) \equiv \sum_{i \in I} d_i$. For $a \neq b$ (due to (4.15)) we obtain

$$d(I_s \cap I_l) = \frac{d(I_s) d(I_l)}{D - 2}, \tag{5.3}$$

for all $s \in S_a$, $l \in S_b$, while for $a = b$ we get (see (2.16))

$$2 \left[d(I_s \cap I_l) - \frac{d(I_s) d(I_l)}{D - 2} \right] = A_{sl} \left[d(I_l) - \frac{(d(I_l))^2}{D - 2} \right], \tag{5.4}$$

$s, l \in S_a$.

Thus the intersection rules do not depend upon parameters $q_{(a)} > 0$. They depend upon the choice of quasi-Cartan matrix A_{sl} (e.g. Cartan one), which has a block-diagonal structure due to (4.15).

q-analogues of $M2 \cap M2$, $M2 \cap M5$ and $M5 \cap M5$ solutions corresponding to the semisimple Lie algebra $\mathbf{A_1} \oplus \mathbf{A_1}$ were suggested in [4].

In [5] a simulation of $M_2 \cap M_5$ dyonic configuration in $D = 11$ supergravity [21] was considered. It corresponds to the Lie algebra $\mathbf{A_2}$. Here we consider a q-extension of this solution with $q = 1, 2, \ldots$.

The solution is defined on the manifold

$$M = (2\mu, +\infty) \times (M_0 = S^2) \times (M_1 = \mathbb{R}) \times M_2 \times M_3, \tag{5.5}$$

where $\dim M_2 = 2$ and $\dim M_3 = 5$. The U^s-vectors corresponding to fluid components obey (5.1) with $I_1 = \{1, 2\}$, $I_2 = \{1, 3\}$ and $q_{(a)} = q$.

The solution reads as following

$$g = H_1^{1/(3q)} H_2^{2/(3q)} \Big\{ F^{-1} dR \otimes dR + R^2 h[S^2] - F H_1^{-1/q} H_2^{-1/q} dt \otimes dt \tag{5.6}$$

$$+ H_1^{-1/q} h^{[2]} + H_2^{-1/q} h^{[3]} \Big\},$$

$$\rho^1 = - \frac{(2q-1) A_1 F^{q-1}}{J_0 R^4} H_1^{-2/q} H_2^{1/q}, \tag{5.7}$$

$$\rho^2 = - \frac{(2q-1) A_2 F^{q-1}}{J_0 R^4} H_1^{1/q} H_2^{-2/q}, \tag{5.8}$$

where $F = 1 - 2\mu/R$, $J_0 = H_1^{1/(3q)} H_2^{2/(3q)}$; $h[S^2]$ is the canonical metric on 2-dimensional sphere S^2, $h^{[2]}$ and $h^{[3]}$ are Ricci-flat metrics of Euclidean signatures defined on the manifolds M_2 and M_3, respectively; $\mu > 0$ and $H_s(Z)$ are defined in (4.12), with $Z = Z(z, q)$ from (4.3), $z = R^{-1}$ and parameters P_s, $P_s^{(2)}$, $\bar{B}_s = 4A_s$, $s = 1, 2$ obey (4.13). Here $K_s = (U^s, U^s) = 2q^2$ for $s = 1, 2$.

This solution for $q = 1$ simulates $\mathbf{A_2}$-dyon from [21] consisting of an electric $M2$-brane with a worldvolume isomorphic to $(M_1 = \mathbb{R}) \times M_2$ and a magnetic $M5$-brane with a worldvolume isomorphic to $(M_1 = \mathbb{R}) \times M_3$ [5]. The branes are intersecting on the time manifold $M_1 = \mathbb{R}$.

Hawking temperature.

The Hawking temperature corresponding to a solution with a horizon (3.1) has the following form (see [34])

$$T_H = \frac{d}{4\pi (2\mu)^{1/d}} \prod_{s=1}^{m} H_{s0}^{-q_s h_s}, \tag{5.9}$$

where H_{s0} are defined in (3.8).

For the q-extension of dyonic solution from the previous subsection we get

$$T_H = \frac{1}{8\pi\mu} (H_{10} H_{20})^{-1/(2q)}. \tag{5.10}$$

Here

$$H_{s0} = 1 + P_s (2\mu)^{-1} + P_s^{(2)} (2\mu)^{-2}, \tag{5.11}$$

where $\mu > 0$ and parameters $P_s > 0$, $P_s^{(2)} > 0$ obey (4.13) with $\bar{B}_s = 4A_s$, $A_s < 0$.

For fixed parameters $\mu > 0$ and $P_s > 0$ (and hence $A_s < 0$) the values $H_{s0} > 1$ do not depend upon q. The Hawking temperature $T_H(q)$ is monotonically increasing in a sequence $q = 1, 2, \dots$ and tends to the Schwarzschild value $T_H(+\infty) = 1/(8\pi\mu)$ as $q \to +\infty$. In this limit the metric (5.7) tends to $g_{Sch} + h^{[2]} + h^{[3]}$, where g_{Sch} is the Schwarzschild metric. The densities ρ^s and pressures p_i^s vanish as $q \to +\infty$.

For fixed parameters $P_s > 0$ and q the Hawking temperature has the following asymptotical behaviour $T_H(\mu) \sim \mathrm{const}\mu^{2/q-1}$ as $\mu \to +0$. It is divergent for $q > 2$, i. e. $T_H(+0) = +\infty$. For $q = 1$ we get $T_H(+0) = 0$ while for $q = 2$ we find $T_H(+0) = T_0 > 0$.

We note that the analysis of T_H may be of relevance for possible search of dual holographic models in a context of (an extended version of) AdS/CFT approach [35, 36, 37].

6 Conclusions

In this paper we have considered a family of spherically symmetric solutions in the model with m-component anisotropic fluid when the equations of state (1.1)- (1.2) (with certain restrictions on U_i^s-parameters) were imposed. The metric of any solution contains $(n-1)$ Ricci-flat “internal” space metrics and depends upon a set of parameters U_i^s which define the the equations of state of fluid components. The solutions under consideration are governed by moduli functions H_s obeying non-linear differential equations with certain boundary conditions imposed.

The parameters $U_1^s = q_s > 0$ play a special role among all other parameters U_i^s. Here we have shown that for coinciding $q_s = q$ there exists a subclass of solutions with regular horizon when $q = 1, 2, \ldots$ and U^s-vectors correspond to simple roots of certain semisimple Lie algebra. The regularity of horizon follow from the polynomial structure of the functions $H_s(Z)$ and $Z = Z(z)$ (see (4.3)).

We have also generalized one-block solutions with regular horizon to the case of block-orthogonal set of vectors U^s. In this case several natural parameters $q_{(a)}$ occur, $a = 1, \ldots, k$, where k is a number of blocks. We have also suggested generalized analogues of black brane solutions in the block-orthogonal case.

As an example we have presented a q-analogue of $M_2 \cap M_5$ dyonic solution in $D = 11$ supergravity (corresponding to Lie algebra A_2) with $q = 1, 2, \ldots$. We have shown that for a fixed parameter of extremality $\mu > 0$ and fluid parameters $P_s > 0$ the Hawking temperature T_H is monotonically increasing in a sequence $q = 1, 2, \ldots$ and tends to the Schwarzschild value $1/(8\pi\mu)$ as $q \to +\infty$. Here an open problem is a search of dual holographic quantum models (with temperatures) corresponding to these (classical) solutions.

Acknowledgments

This work was supported in part by Russian Foundation for Basic Research (Grant Nr. 09-02-00677-a) and by the FTsP "Nauchnie i nauchno-pedagogicheskie kadry innovatsionnoy Rossii" for the years 2009-2013.

References

[1] V.D. Ivashchuk, V.N. Melnikov and A.B. Selivanov, Multidimensional black hole solutions in model with anisotropic fluid, *Grav. Cosmol.* **7**, 4(12), 308-310 (2001); gr-qc/0205103.

[2] V.D. Ivashchuk, V.N. Melnikov and A.B. Selivanov, Simulation of intersecting black brane solutions by multicomponent anisotropic fluid, *Grav. Cosmol.* **9**, No. 1-2 (33-34), 50-54 (2003); hep-th/0211247.

[3] H. Dehnen, V.D. Ivashchuk and V.N. Melnikov, On black hole solutions in model with anisotropic fluid, *Grav. Cosmol.* **9**, No. 3 (35), 153-158 (2003); gr-qc/0211049.

[4] H. Dehnen and V.D. Ivashchuk, On spherically symmetric solutions with horizon in model with multicomponent anisotropic fluid, *J. Math. Phys.* **45**, 4726-4736 (2004); gr-qc/0310043.

[5] V.D. Ivashchuk On analogues of black brane solutions in the model with multicomponent anisotropic fluid, *Phys. Lett.* **B 693**, No. 3,4, (2010) 399403; arXiv: 1001.4053.

[6] K.S. Stelle, Lectures on supergravity p-branes, hep-th/9701088.

[7] G.T. Horowitz and A. Strominger, Black string and p-branes, *Nucl. Phys.* **B 360**, 197-209 (1991)

[8] M.J. Duff and J.X. Lu, Elementary 5-brane solutions of D=10 supergravity, *Nucl. Phys.* **B 354**, 141-153 (1991).

[9] M. Cvetic and A. Tseytlin, Non-extreme black holes from non-extreme intersecting M-branes, *Nucl. Phys.* **B 478**, 181-198 (1996); hep-th/9606033.

[10] H. Lü and C.N. Pope, Black p-branes and their vertical dimensional reduction, *Nucl. Phys.* **B 489**, 264-278 (1997); hep-th/9609126.

[11] M. Costa, Black composite M-branes,*Nucl. Phys.* **B 490**, 603-614 (1997); hep-th/9610138.

[12] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-extremal intersecting p-branes in various dimensions, *Phys. Lett.* **B 406**, 44-48 (1997); hep-th/9702079.

[13] N. Ohta, Intersection rules for non-extreme p-branes, *Phys. Lett.* **B 403**, 218-224 (1997); hep-th/9702164.

[14] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, The Reissner-Nordström problem for intersecting electric and magnetic p-branes, *Grav. Cosmol.* **3**, 3(11), 203-212 (1997); gr-qc/9710054.

[15] H. Lü, J. Maharana, S. Mukherji and C.N. Pope, Cosmological solutions, p-branes and the Wheeler-DeWitt equation, *Phys. Rev.* **D 57**, 2219-2229 (1997); hep-th/9707182.

[16] V.D. Ivashchuk and V.N. Melnikov, Multidimensional classical and quantum cosmology with intersecting p-branes, *J. Math. Phys.*, **39**, 2866-2889 (1998); hep-th/9708157.

[17] K.A. Bronnikov, Block-orthogonal brane systems, black holes and wormholes, *Grav. Cosmol.* **4**, No 1 (13), 49 (1998); hep-th/9710207.

[18] V.D. Ivashchuk and V.N. Melnikov, In Lecture Notes in Physics, Vol. 537, "Mathematical and Quantum Aspects of Relativity and Cosmology Proceedings of the Second Samos Meeting on Cosmology, Geometry and Relativity held at Pythagoreon, Samos, Greece, 1998, eds: S. Cotsakis, G.W. Gibbons., Berlin, Springer, pp. 214-247, 2000; gr-qc/9901001.

[19] S. Cotsakis, V.D. Ivashchuk and V.N. Melnikov. P-brane Black Holes and Post-Newtonian Approximation, *Grav. Cosmol.*, **5**, No 1 (17), 52-57 (1999); gr-qc/9804080.

[20] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.*, **41** (1) 444-460 (2000); hep-th/9907019.

[21] V.D. Ivashchuk and V.N. Melnikov, P-brane black holes for general intersections, *Grav. Cosmol.*, **5**, No 4 (20) 313-318 (1999); gr-qc/0002085.

[22] V.D. Ivashchuk and V.N. Melnikov, Black hole p-brane solutions for general intersection rules, *Grav. Cosmol.*, **6**, No 1 (21) 27-40 (2000); hep-th/9910041.

[23] V.D. Ivashchuk and V.N. Melnikov, Toda p-brane black holes and polynomials related to Lie algebras, *Class. Quantum Grav.*, **17** 2073-2092 (2000); math-ph/0002048.

[24] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.*, **18** (2001) R87-R152; hep-th/0110274.

[25] M.A. Grebeniuk, V.D. Ivashchuk and S.-W. Kim, Black-brane solutions for C_2 algebra, *J. Math. Phys.* **43**, 6016-6023 (2002); hep-th/0111219.

[26] M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov Black-brane solution for A_3 algebra, *Phys. Lett.* , **B 543**, 98-106 (2002); hep-th/0208083.

[27] V.D. Ivashchuk and V.N. Melnikov, On black brane solutions and their fluid analogues, *Grav. Cosmol.*, **17**, **4**, 328-334 (2011).

[28] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, On Wheeler-DeWitt equation in multidimensional cosmology, *Nuovo Cimento* **B 104**, 5, 575-581 (1989).

[29] V.D. Ivashchuk and V.N. Melnikov, Multidimensional cosmology with m-component perfect fluid, *Int. J. Mod. Phys.* **D 3**, 4, 795-811 (1994); gr-qc/ 9403063.

[30] D.E. Liebscher and U. Bleyer, Multifactor cosmological models, *Grav. Cosmol.*, **1**, No 1 (1), 31-36 (1995).

[31] V.R. Gavrilov, V.D. Ivashchuk and V.N. Melnikov, Integrable pseudo-Euclidean Toda-like systems in multidimensional cosmology with multicomponent perfect fluid, *J. Math. Phys* **36**, 5829-5847 (1995).

[32] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and Representations. A graduate course for physicists, Cambridge University Press, Cambridge, 1997.

[33] V.D. Ivashchuk, in preparation.

[34] J.W. York, *Phys. Rev.* **D 31**, 775 (1985).

[35] J.M. Maldacena, *Adv. in Theor. Math. Phys.* **2**, 231-252 (1998); hep-th/9711200.

[36] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Phys. Lett.* **B 428**, 105-114 (1998); hep-th/9802109.

[37] E. Witten (1998), *Adv. in Theor. Math. Phys.* **2**, 253-291 (1998); hep-th/9802150.